arXiv:1412.5450v2 [math.AG] 1 Jul 2015

A BOTT TYPE RESIDUE FORMULA ON COMPLEX ORBIFOLDS

MAURÍCIO CORRÊA JR., MIGUEL RODRIGUEZ PEÑA AND MÁRCIO G. SOARES

ABSTRACT. We prove residual formulas for vector fields defined on compact complex orbifolds with isolated singularities and give some applications of these on weighted projective spaces.

To José Seade on the occasion of his 60th birthday

INTRODUCTION

In this article we prove residual formulas for vector fields defined on compact complex orbifolds with isolated singularities. We then derive some applications, related to holomorphic foliations on weighted complex projective spaces and conclude by looking at Hirzebruch surfaces as resolutions of weighted projective planes of a simple type. These residual formulas are inspired by the well known residue formula of R. Bott for vector fields, with isolated zeros, defined on complex manifolds and also by several more recent results on this subject, which can be found in [3].

V-manifolds were introduced by I. Satake in [11] and later rediscovered by W. Thurston [12], which referred to them as *orbifolds* and since then they are known by this name. These objects may present singularities, but these are all quotient singularities originating locally from the action of finite groups and hence are tractable. More precisely, an orbifold X is a complex space endowed with the following property: each point $p \in X$ possesses a neighborhood which is the quotient Y/G_p, where Y is a complex manifold, say of dimension n, and G_p is a properly discontinuous finite group of automorphisms of Y, so that locally we have a quotient map $(Y, \tilde{p}) \xrightarrow{\pi_p} (Y/G_p, p) = (X, p)$. If the isotropy group G_p of p is non-trivial, then p is a singular point of X. The structure of orbifolds around its singularities was elucidated by D. Prill in [10]. We will also make use of results by R. Blache [2] concerning Chern classes and local Chern classes on orbifolds. An extensive account on orbifolds can be found in [1].

In Section 2 we state and prove the main results of this note, which we briefly present now.

Theorem 1 *Let X be a compact orbifold of dimension n with only isolated singularities, let $\mathscr{L}$ be a locally V-free sheaf of rank 1 over X and L the associated line V-bundle. Suppose ξ is a holomorphic section of $TX \otimes L$ with isolated zeros. For each zero p of the section ξ, let $\tilde{\xi} = \pi_p^* \xi$ be a local lifting of ξ via the quotient map $(Y, \tilde{p}) \xrightarrow{\pi_p} (X, p)$ and let $J\tilde{\xi} = \left(\dfrac{\partial \tilde{\xi}_i}{\partial \tilde{z}_j} \right)_{1 \leq i,j \leq n}$ be the matrix of the linear part of $\tilde{\xi}$. If P is any invariant*

[1] *2000 Mathematics Subject Classification.* 57R18, 55N32

[2] *Key words:* orbifolds, Chern classes, indices of vector fields.

[3] *This work was partially supported by CNPq, CAPES, FAPEMIG and FAPERJ.*

polynomial of degree n and $L^\vee$ is the dual of L then

$$\int_X P(TX - L^\vee) = \sum_{p\,|\,\xi(p)=0} \frac{1}{\#G_p}\mathrm{Res}_{\tilde{p}}\left[\frac{P(J\tilde{\xi})\, d\tilde{z}_1 \wedge \cdots \wedge d\tilde{z}_n}{\tilde{\xi}_1 \ldots \tilde{\xi}_n}\right]$$

where $Res_{\tilde{p}}\left[\dfrac{P(J\tilde{\xi})\, d\tilde{z}_1 \wedge \cdots \wedge d\tilde{z}_n}{\tilde{\xi}_1 \ldots \tilde{\xi}_n}\right]$ is Grothendieck's point residue.

Theorem 2 *Let X be a compact complex orbifold of dimension n with only isolated singularities, $\mathscr{L}$ a locally V-free sheaf of rank 1 over X and L the associated line V-bundle. Let $\sigma : \widetilde{X} \longrightarrow X$ be a good resolution of X, $\mathcal{E}$ the exceptional divisor, and let ξ be a holomorphic section of $TX \otimes L$ with isolated zeros. Assume $(\sigma^*\mathscr{L})^{\vee\vee}$ is locally free. Then*

$$\sum_{p\,|\,\xi(p)=0} I_p(\xi) = \int_{\widetilde{X}} c_n\left((\sigma^*(TX \otimes L))^{\vee\vee}\right) - \sum_{p\in Sing(X)} \int_{\widetilde{X}} c_n\left(p, (\sigma^*(TX \otimes L))^{\vee\vee}\right).$$

In particular, if $\sigma^*(\mathscr{T}_X \otimes \mathscr{L})$ is reflexive, then

$$\sum_{p\in Sing(X)\cap\{\xi(p)=0\}} I_p(\xi) = \sum_{\tilde{p}\in\mathcal{E}\,|\,\sigma^*\xi(\tilde{p})=0} I_{\tilde{p}}(\sigma^*\xi) - \sum_{p\in Sing(X)} \int_{\widetilde{X}} c_n\left(p, \sigma^*(TX \otimes L)\right),$$

where $I_p(\xi)$ is the Poincaré-Hopf index of the section ξ at p.

In Section 3 we exploit some consequences of these results in weighted projective spaces $\mathbb{P}^n_w$, among which an enumerative result counting the number of singularities of holomorphic foliations on these spaces and also, in a particular but interesting case, a condition for a holomorphic foliation on $\mathbb{P}(1,1,k)$ to have a singularity at the singular point $\mathsf{e}_2 = [0 : 0 : 1]_{\mathsf{k}}$ of $\mathbb{P}(1,1,k)$. The latter is shown in two different ways, the first one is based on the enumerative result for $\mathbb{P}^n_w$ mentioned above, and the second one is obtained by considering the Hirzebruch surfaces Σ_k as good resolutions of $\mathbb{P}(1,1,k)$. It's worth pointing out that if a holomorphic foliation $\mathcal{F}_{|\mathbb{P}(1,1,k)\setminus\{\mathsf{e}_2\}}$ is given, around the singular point $\mathsf{e}_2 \in \mathbb{P}(1,1,\mathsf{k})$, by a vector field ξ or by a differential 1-form η, then the lifted vector field $\tilde{\xi}$ or the lifted 1-form $\tilde{\eta}$ on $(\mathbb{C}^3,0)$, via $(\mathbb{C}^3,0) \xrightarrow{\pi_{\mathsf{e}_2}} (\mathbb{C}^3/G_{\mathsf{e}_2}, \mathsf{e}_2)$, extend holomorphically to 0 and this extension may be non-vanishing at 0. If this is the case we say that $\mathcal{F}$ is non-singular at e_2 (see Example 3.10).

1. Preliminaries

1.1. **Orbifolds or V-manifolds.** We will make use of the theory of Chern classes for pairs $(X, \mathscr{S})$, as developed by R. Blache in [2], where X is a complex orbifold with isolated quotient singularities and $\mathscr{S}$ is a coherent sheaf on X. In fact, what we will exploit is a little more restricted, namely, the Chern classes for locally free V-sheaves on orbifolds. So, we start by briefly recalling the essential objects we will consider. We refer the reader to [1], [2] and [11] for the definitions and results on orbifolds that we present in this subsection.

Definition 1.1. *An orbifold is a connected paracompact complex space X satisfying the property that each point $x \in X$ has an open neighborhood $U \subset X$ which is a quotient $U \cong V/G$, where V is a complex manifold, G is a finite subgroup of the group of biholomorphisms of V, with V and G depending on x. (X, x) denotes the germ of the orbifold*

X at the point x and is referred to as a quotient germ. Morphisms of orbifolds are the holomorphic maps between them.

Since the possible singularities appearing in a orbifold X are quotient singularities, we have that X is reduced, normal, Cohen-Macaulay and with only rational singularities. We denote its singular set by $Sing(X)$ and the regular part of X by $X_{reg} = X \setminus Sing(X)$. The dimension of X is the dimension of the complex manifold X_{reg}.

The local structure of orbifolds around its singularities was described by D. Prill in [10]: if $\dim X = n$, then every quotient germ (X, x) determines a unique (up to conjugation) small finite group $G_x \subset Gl(n, \mathbb{C})$ such that $(X, x) = (Y, y)/G_x$, where $(Y, y) = (\mathbb{C}^n, 0)$, with natural projection $\pi : (Y, y) \to (X, x)$ which is called the *local smoothing covering* of X at x. Note that $x \in Sing(X)$ precisely when the isotropy group G_x is non-trivial. We point out that $G_x \cong \pi_1((X, x) \setminus Sing(X))$ and for this reason G_x is called the *local fundamental group* of X at x. Recall that G_x is *small* if $\operatorname{codim}_{\mathbb{C}} \operatorname{Fix} G_x \geq 2$, or equivalently, if no element of G_x has 1 as an eigenvalue of multiplicity $n - 1$.

Satake's fundamental idea was to use local smoothing coverings to extend the definition of usual known objects to orbifolds. For instance, smooth differential k-forms on X are C^∞-differential k-forms ω on X_{reg} such that the pull-back $\pi^*\omega$ extends to a C^∞-differential k-form on every local smoothing covering $\pi : (Y, y) \to (X, x)$ of X. Hence, if ω is a smooth $2n$-form on X with compact support $\operatorname{Supp}(\omega) \subset (X, x)$ then, by definition,

$$\int_X \omega = \frac{1}{\#G_x} \int_Y \pi^*\omega. \tag{1}$$

If now the $2n$-form ω has compact support, we use a partition of unity $\{\rho_\alpha, U_\alpha\}_{\alpha \in \mathcal{A}}$, where $\pi_\alpha : (\widetilde{U}_\alpha, \tilde{p}_\alpha) \longrightarrow (U_\alpha, p_\alpha)$ is a local smoothing covering and $\sum \rho_\alpha(x) = 1$ for all $x \in \operatorname{Supp}(\omega)$, and set

$$\int_X \omega = \sum_\alpha \int_X \rho_\alpha \omega. \tag{2}$$

Analogously to the manifold case we have the sheaf $\mathscr{E}^k$ of C^∞-differential k-forms on X, the exterior differentiation $d : \mathscr{E}^k \to \mathscr{E}^{k+1}$ and $\mathscr{E}^* = \bigoplus_{0 \leq k \leq 2n} \mathscr{E}^k$.

Stokes' formula also holds, in particular, if X has dimension n and $\omega \in \Gamma_c(X, \mathscr{E}^{2n-1})$ (compact supports) then

$$\int_X d\omega = 0. \tag{3}$$

The de Rham complexes $H^*_{dR}(X, \mathbb{C}) = \bigoplus_{0 \leq k \leq 2n} H^k_{dR}(X, \mathbb{C})$ and with compact support $H^*_{dRc}(X, \mathbb{C}) = \bigoplus_{0 \leq k \leq 2n} H^k_{dRc}(X, \mathbb{C})$ are then well defined and Poincaré's duality holds:

the pairing

$$\begin{array}{ccc} H^k_{dR}(X,\mathbb{C}) \times H^{2n-k}_{dRc}(X,\mathbb{C}) & \longrightarrow & \mathbb{C} \\ ([\omega],[\eta]) & \longmapsto & \displaystyle\int_X \omega \wedge \eta \end{array} \tag{4}$$

is nondegenerate. As in the case of manifolds, de Rham's theorem holds on orbifolds and we have $H^*(X,\mathbb{C}) \cong H^*_{dR}(X,\mathbb{C})$, where $H^*(X,\mathbb{C})$ is the complex of singular cohomology rings with complex coefficients and, by considering Čech cohomology we also get $H^*_{dR}(X,\mathbb{C}) \cong \check{H}^*(X,\mathbb{C})$.

By invoking Hironaka's resolution of singularities we conclude that, given $x \in X$, there exists an open neighborhood $U \subset X$ of x such that, if $\sigma : \widetilde{U} \longrightarrow U$ is a resolution, $\widetilde{Y}$ is a resolution of the normalization of the fiber product $\widetilde{U} \times_U Y$, where $\pi : Y \longrightarrow U$ is a local smoothing covering, then there is a commutative diagram

$$\begin{array}{ccc} \widetilde{Y} & \xrightarrow{\phi} & Y \\ \psi \downarrow & & \downarrow \pi \\ \widetilde{U} & \xrightarrow{\sigma} & U \end{array} \tag{5}$$

with ϕ bimeromorphic and biholomorphic over $\pi^{-1}(U \setminus Sing(U))$ and ψ generically finite of $\deg \psi = \deg \pi$ and unramified over $\widetilde{U} \setminus \sigma^{-1}(Sing(U))$.

Definition 1.2. *Given an irreducible normal complex space Y and $W \subsetneq Y$ a connected compact analytic subset, let (Y,W) denote the germ of Y along W. A connected open subset $V \subset Y$ is a good representative of (Y,W) if $W \subset V$, ∂V is a real C^∞-manifold and there is a contraction of V onto W. If V is a good representative of (Y,W) we define $H^*_{dRc}((Y,W)) := H^*_{dRc}(V,\mathbb{C})$.*

The definition is good since $H^*_{dRc}(V) = H^*_{dRc}(V')$ if V and V' are good representatives of (Y,W). Note that if both Y and W are manifolds with $\mathrm{codim}_{\mathbb{C}} W = 1$, then $H^k_{dRc}((Y,W)) \cong H^{k-2}_{dR}(W)$ for $k \geq 2$.

Definition 1.3. *A good resolution of an orbifold X is a resolution $\sigma : \widetilde{X} \longrightarrow X$ whose exceptional divisor $\mathcal{E}$ has only normal crossings. If X has only isolated singularities we also require that $\sigma(\mathcal{E}) = Sing(X)$.*

Good resolutions always exist.

1.2. **Locally V-free sheaves, V-bundles and Chern classes.** We recall the concept of a coherent G-sheaf on a complex space:

Definition 1.4. *Let $\mathscr{S}$ be a coherent sheaf on a complex space X, let G be a group and $G \longrightarrow Aut(X)$ be a group homomorphism. G acts on $\mathscr{S}$ over X provided:*

(i) *given an open subset $U \subseteq X$ such that $g(U) = U$ for all $g \in G$, G acts on the $\mathbb{C}$-linear space $\Gamma(U,\mathscr{S})$*
(ii) *these actions are compatible with all restriction maps induced by open inclusions $U' \hookrightarrow U$,*
(iii) *$g(f \cdot \xi) = g^* f \cdot g(\xi)$ for all $\xi \in \Gamma(U,\mathscr{S})$, $f \in \Gamma(U,\mathscr{O}_U)$, $g \in G$.*

In case G acts properly discontinuously on X and $\mathscr{S}$ is a coherent G-sheaf on X then, for all $x \in X$ and for all $g \in G$, the action on X induces an identification $\mathscr{O}_{X,x} \simeq \mathscr{O}_{X,g\cdot x}$ and analogously $\mathscr{S}_x \simeq \mathscr{S}_{g\cdot x}$ by the action on $\mathscr{S}$. In particular, if $g \cdot x = x$ for all g, then G acts on the stalk $\mathscr{S}_x$.

Definition 1.5. *Let (X, x) be a quotient germ and $\pi : (Y, y) \longrightarrow (X, x)$ be a local smoothing covering. A coherent sheaf $\mathscr{S}$ on (X, x) is V-free if the following equivalent conditions are satisfied:*

(i) *$\mathscr{S}$ is reflexive (that is, $\mathscr{S}$ is isomorphic to $\mathscr{S}^{\vee\vee}$) and $(\pi^*\mathscr{S})^{\vee\vee}$ is a free sheaf.*
(ii) *There is a free sheaf $\breve{\mathscr{S}}$ on (Y, y) such that G_x acts on $\breve{\mathscr{S}}$ and $\mathscr{S} \simeq \pi_*^{G_x}(\breve{\mathscr{S}})$, where $\pi_*^{G_x}(\breve{\mathscr{S}})$ is the maximal subsheaf of $\pi_*(\breve{\mathscr{S}})$ on which G_x acts trivially.*

A coherent sheaf $\mathscr{S}$ on an orbifold X is locally V-free if $\mathscr{S}_{|(X,x)}$ is V-free for all $x \in X$.

Remark that $\breve{\mathscr{S}} = (\pi^*\mathscr{S})^{\vee\vee}$.

Definition 1.6. *A V-bundle E on an orbifold X is a holomorphic vector bundle E_{reg} on X_{reg} such that $\mathsf{i}_*(\mathscr{O}(E_{reg}))$ is a locally V-free sheaf. Here $\mathsf{i} : X_{reg} \hookrightarrow X$ is the inclusion and $\mathscr{O}(E_{reg})$ is the sheaf of sections of E_{reg}.*

Observe that a holomorphic vector bundle E_{reg} on X_{reg} is a V-bundle on X if, and only if, for every local smoothing covering $\pi : (Y, y) \longrightarrow (X, x)$, there is a holomorphic vector bundle $\breve{E}_Y$ on (Y, y) and an action of G_x on $\breve{E}_Y$ with

$$\left((\breve{E}_Y)_{|(Y,y)\setminus \operatorname{Fix} G_x}\right)/G_x \simeq E_{reg|(X,x)}.$$

Also, $\breve{E}_Y$ and the G_x action on $\breve{E}_Y$ are uniquely determined by E_{reg}.

Now, locally V-free sheaves have the following property: if $\mathscr{S}$ is a locally V-free sheaf on X, then we can associate to it a V-bundle E on X (unique up to isomorphism) satisfying: $\mathscr{S}$ is isomorphic to the sheaf of sections of E.

To construct Chern classes we will use the Chern-Weil procedure via metric connections. First we need to define metrics.

Definition 1.7. *Given an orbifold X and a locally V-free sheaf $\mathscr{S}$ on X, let E be the V-bundle associated to $\mathscr{S}$. A V-metric h on $\mathscr{S}$ is a hermitian metric on E_{reg} (the restriction of E to X_{reg}) such that π^*h admits an extension to a hermitian metric $\breve{h}$ on $\breve{E}_Y$ for all local smoothing coverings $\pi : (Y, y) \longrightarrow (X, x)$.*

As in the case of manifolds, by using partitions of unity we can construct V-metrics.

Given an orbifold X of dimension n, a locally V-free sheaf $\mathscr{S}$ of rank r and a V-metric h on $\mathscr{S}$ we construct classes $c_i(X, \mathscr{S}, h) \in H^{2i}_{dR}(X, \mathbb{C})$, $i = 0, \ldots, n$, as follows:

On the regular part X_{reg} take the metric connection ∇_h on $\mathscr{S}$ associated to h and its curvature operator $K_{\nabla_h} = \nabla_h \circ \nabla_h$, which is represented by a $r \times r$ matrix of smooth 2-forms, say Θ_h, in any local trivialization of $\mathscr{S}$. If $M(r, \mathbb{C})$ is the algebra of $r \times r$ matrices over $\mathbb{C}$, let $C_i : M(r, \mathbb{C}) \longrightarrow \mathbb{C}$ be the i-th elementary symmetric invariant polynomial and define the Chern form $c_i(h) := C_i\left(\frac{\sqrt{-1}}{2\pi}\Theta_h\right) \in \Gamma(X_{reg}, \mathscr{E}^{2i})$. Doing this on all local smoothing coverings we conclude that in fact $c_i(h) \in \Gamma(X, \mathscr{E}^{2i})$. It follows, as in the C^∞ case, that $d(c_i(h)) = 0$ for all i and that $c_i(\mathscr{S}) = [c_i(h)] \in H^{2i}_{dR}(x, \mathbb{C})$ is independent of the metric.

As usual, the *total Chern class* of $\mathscr{S}$ is defined by $c(\mathscr{S}) := c_0(\mathscr{S}) + \cdots + c_r(\mathscr{S}) + \cdots + c_n(\mathscr{S})$, with $c_0(\mathscr{S}) := 1$.

If $\mathscr{T}_X$ is the tangent sheaf of X (which is reflexive), then the *Chern classes* of X are, by definition, $c_i(X) = c_i(\mathscr{T}_X)$, $i = 0, \ldots, n$. Remark that $\mathscr{T}_X$ is locally V-free and $\mathscr{T}_{X|X_{reg}}$ is the holomorphic tangent sheaf of the complex manifold X_{reg}.

The usual properties of the Chern classes hold with the following adjustments: a sequence $\mathscr{S}_m \longrightarrow \cdots \longrightarrow \mathscr{S}_1$ of locally V-free sheaves on X is *V-exact* provided the sequence

$$(\pi^*\mathscr{S}_m)^{\vee\vee} \longrightarrow \cdots \longrightarrow (\pi^*\mathscr{S}_1)^{\vee\vee}$$

is exact, for all local smoothing coverings of X. With this in mind we have:

Remark 1. *Properties of the Chern classes.*

(i) *if $\mathscr{S}$ is locally V-free and has rank r, then $c_i(\mathscr{S}) = 0$ for $i > r$.*
(ii) $c_i(\mathscr{S}) = (-1)^i\, c_i(\mathscr{S}^\vee)$ *for* $i \geq 0$.
(iii) $c(\mathscr{S}) = c(\mathscr{S}')\, c(\mathscr{S}'')$ *whenever we have a V-exact sequence of locally V-free sheaves* $0 \longrightarrow c(\mathscr{S}') \longrightarrow c(\mathscr{S}) \longrightarrow c(\mathscr{S}'') \longrightarrow 0$.
(iv) $c_i((\mathscr{S} \otimes \mathscr{L})^{\vee\vee}) = \sum_{k=0}^{i} \binom{r-k}{i-k} c_k(\mathscr{S})\, c_1(\mathscr{L})^{i-k}$ *for* $0 \leq i \leq r$, *where $\mathscr{L}$ is any locally V-free sheaf of rank* 1 *on* X.
(v) $c(\mathscr{S}) \in H^*(X, \mathbb{Q})$.

Property *(v)* above is due to the natural use of local smoothing coverings which implies, in many situations, that we divide by the orders of the groups involved. It follows by showing that

$$\xi^r + \xi^{r-1} p^* c_1(\mathscr{S}) + \cdots + \xi p^* c_{r-1}(\mathscr{S}) + p^* c_r(\mathscr{S}) = 0$$

in $H^*(\mathbb{P}(\mathscr{S}), \mathbb{Q})$, where $\mathbb{P}(\mathscr{S}) \xrightarrow{p} X$ is the projectivization of $\mathscr{S}$ over X, with $\xi \in H^2(\mathbb{P}(\mathscr{S}), \mathbb{Q})$ given by $c_1(\mathscr{O}_{\mathbb{P}(\mathscr{S})}(1))$ and $c_i(\mathscr{S}) \in H^{2i}(X, \mathbb{Q})$ uniquely determined.

Chern numbers are defined as in the smooth case, that is, suppose X is a compact orbifold of dimension n and let $\mathscr{S}$ be a locally V-free sheaf on X. For $\mathbb{K} = \mathbb{Q}$ or $\mathbb{C}$ the map

$$\begin{array}{ccc} H^{2n}_{dR}(X, \mathbb{K}) & \longrightarrow & \mathbb{K} \\ [\omega] & \longmapsto & [X] \frown [\omega] \end{array} \tag{6}$$

gives an identification $H^{2n}_{dR}(X, \mathbb{K}) \simeq \mathbb{K}$. In $\mathbb{K}[Z_1, \ldots, Z_n]$ give weight (or degree) i to Z_i. If P is a quasihomogeneous monomial of degree n, $P = Z_1^{\nu_1} \ldots Z_n^{\nu_n}$, $\nu_1 + 2\nu_2 + \cdots + n\nu_n = n$, then the number $P(c(\mathscr{S})) := c_1(\mathscr{S})^{\nu_1} \cdots c_n(\mathscr{S})^{\nu_n} \in \mathbb{K}$, obtained via (6), is the *Chern number of $\mathscr{S}$ with respect to* $(\nu_1, \ldots, \nu_n)$. If $P \in \mathbb{K}[Z_1, \ldots, Z_n]$ is a quasihomogeneous polynomial of degree n, then the number $P(c(\mathscr{S})) \in \mathbb{K}$ is defined in the same way.

1.3. **Local Chern classes.** R. Blache [2] developed a natural concept of local Chern classes on orbifolds, which will be very useful for our purposes. We briefly describe it.

Let $\sigma : (\widetilde{X}, \mathcal{E}) \longrightarrow (X, x)$ be a good resolution of an isolated quotient singularity as in Definition 1.3 and suppose $\breve{\mathscr{S}}$ is a locally free sheaf on $\widetilde{X}$ such that $\mathscr{S} = (\sigma_*\breve{\mathscr{S}})^{\vee\vee}$ is a V-free sheaf on X. Let $U \subset W$ be open neighborhoods of x with U relatively compact and such that U and W are good representatives of (X, x) as in Definition 1.2. Then:

(i) There exist metrics h and $\breve{h}$ on $\mathscr{S}$ and $\breve{\mathscr{S}}$, respectively, such that $h_{|W\setminus U} = \breve{h}_{|\widetilde{W}\setminus\widetilde{U}}$.

(ii) Let $c_i(h) \in \Gamma(W, \mathscr{E}^{2i})$ and $c_i(\breve{h}) \in \Gamma(\widetilde{W}, \mathscr{E}^{2i})$ be the Chern forms associated to $(W, \mathscr{S}, h)$ and $(\widetilde{W}, \breve{\mathscr{S}}, \breve{h})$, respectively. Then

$$c_i(x, \breve{\mathscr{S}}) := c_i(\widetilde{X}, \mathcal{E}, \breve{\mathscr{S}}) := c_i(\breve{h}) - \sigma^* c_i(h) \in H^{2i}_{dRc}((\widetilde{X}, \mathcal{E}), \mathbb{C}) \tag{7}$$

does not depend on U, W, h and $\breve{h}$ and is called the *i-th local Chern class of $\breve{\mathscr{S}}$ along $\mathcal{E}$*.

Remark that by (i) the local Chern class $c_i(x, \breve{\mathscr{S}})$ localizes at x. The relation between the local Chern numbers and the Chern numbers is given in Proposition 3.14 of [2] and reads:

Proposition 1.8. *Let $\sigma : \widetilde{X} \longrightarrow X$ be a good resolution of a compact orbifold of dimension n with only isolated singularities and let $\breve{\mathscr{S}}$ be a locally free sheaf on $\widetilde{X}$ with $\mathscr{S} = (\sigma_* \breve{\mathscr{S}})^{\vee\vee}$ a locally V-free sheaf on X. If $P \in \mathbb{C}[Z_1, \dots, Z_n]$ is an invariant polynomial of degree n then,*

$$\int_X P(c(\mathscr{S})) = \int_{\widetilde{X}} P(c(\breve{\mathscr{S}})) - \sum_{x \in Sing(X)} \int_{\widetilde{X}} P(c(x, \breve{\mathscr{S}})). \tag{8}$$

2. Bott's residue formula on orbifolds

2.1. **The residue formula.** Let X be a complex orbifold of dimension n and L a line V-bundle over X and consider the Chern classes

$$c_j(TX - L^\vee) := c_j(X) + c_{j-1}(X)c_1(L) + \cdots + (c_1(L))^j, \quad 1 \le j \le n,$$

$$c^\nu(TX - L^\vee) = c_1^{\nu_1}(TX - L^\vee) \dots c_n^{\nu_n}(TX - L^\vee), \quad \nu = (\nu_1, \dots, \nu_n), \tag{9}$$

$$n = \nu_1 + 2\nu_2 + \cdots + n\nu_n.$$

Write $C^\nu = C_1^{\nu_1} \dots C_n^{\nu_n}$ for the quasihomogeneous invariant polynomial of degree n, where C_i is the i-th elementary symmetric function, and remark that any invariant polynomial P of degree n is a linear combination $P = \sum_\nu a_\nu C^\nu$, $a_\nu \in \mathbb{C}$.

Theorem 2.1. *Let X be a compact orbifold of dimension n with only isolated singularities, let $\mathscr{L}$ be a locally V-free sheaf of rank 1 over X and L the associated line V-bundle. Suppose ξ is a holomorphic section of $TX \otimes L$ with isolated zeros. If P is an invariant polynomial of degree n, then*

$$\int_X P(TX - L^\vee) = \sum_{p \,|\, \xi(p)=0} \frac{1}{\#G_p} Res_{\tilde{p}} \left[\frac{P(J\tilde{\xi})\, d\tilde{z}_1 \wedge \cdots \wedge d\tilde{z}_n}{\tilde{\xi}_1 \dots \tilde{\xi}_n} \right]$$

where, for each p such that $\xi(p) = 0$, $(\widetilde{U}, \tilde{p}) \xrightarrow{\pi_p} (U, p)$ is a smoothing covering of X at p, $\tilde{\xi} = \pi_p^* \xi$, $J\tilde{\xi} = \left(\dfrac{\partial \tilde{\xi}_i}{\partial \tilde{z}_j} \right)_{1 \le i,j \le n}$ and $Res_{\tilde{p}} \left[\dfrac{P(J\tilde{\xi})\, d\tilde{z}_1 \wedge \cdots \wedge d\tilde{z}_n}{\tilde{\xi}_1 \dots \tilde{\xi}_n} \right]$ is Grothendieck's point residue.

Proof. Let $p_1, \dots, p_k$ be the zeros of the section ξ and choose points $p_{k+1}, \dots, p_l$ in X such that:

(i) For all $1 \leq \alpha \leq l$, $\pi_\alpha : (\widetilde{U}_\alpha, \tilde{p}_\alpha) \longrightarrow (U_\alpha, p_\alpha)$ is a smoothing covering, U_α is a good representative of $\{p_\alpha\}$ and $\{U_\alpha\}_{\alpha=1,\dots,l}$ is an open cover of X.
(ii) For each zero p_α of ξ, $p_\alpha \notin U_\beta$ for all $\beta \neq \alpha$.
(iii) If $p_\alpha \neq p_\beta$ are zeros of ξ then $U_\alpha \cap U_\beta = \emptyset$, $U_\alpha \cap Sing(X) = \{p_\alpha\}$ in case $p_\alpha \in Sing(X)$ and $U_\alpha \cap Sing(X) = \emptyset$ otherwise.
(iv) If $p_\alpha \in X_{reg}$ then U_α is a trivializing neighborhood for both TX and L, whereas if $p_\alpha \in Sing(X)$, $U_\alpha \setminus \{p_\alpha\}$ is a trivializing neighborhood for both TX and L.

We follow Chern's arguments [4] for the first part of the proof. The constructions below are all well defined, see [11] §2.

Equip the locally V-free sheaves $\mathscr{T}_X$ and $\mathscr{L}$, the tangent sheaf of X and the sheaf whose associated V-bundle is L, respectively, with hermitian V-metrics h and H.

Locally, using $\pi_p : (\mathbb{C}^n, 0) \cong (\widetilde{U}, \tilde{p}) \longrightarrow (U, p)$, we have $\breve{h} = (\breve{h}_{ij})$ and $\breve{H} = a_{\widetilde{U}} > 0$. The (unique) connection which preserves the hermitian structure in $T\widetilde{U}$ has matrix

$$\omega = \partial\breve{h} \cdot \breve{h}^{-1},\ \omega = (\omega_{ij}),\ \omega_{ij} = \sum_k \Gamma^j_{ik} d\tilde{z}_k.$$

The curvature matrix is

$$K_{\nabla_h} = d\omega - \omega \wedge \omega = \overline{\partial}\omega = (\Omega_{ij}) \quad \text{and} \quad \overline{\partial} K_{\nabla_h} = 0.$$

For $\breve{L}_{\widetilde{U}}$ the curvature form is

$$\Phi = -\partial\overline{\partial} \log a_{\widetilde{U}}, \quad a_{\widetilde{U}} = a_{\widetilde{U}'} |\breve{H}_{\widetilde{U}\widetilde{U}'}|.$$

We have

$$\Phi = -\sum_k d\tilde{z}_k \wedge \psi_k, \ \ \psi_k = \overline{\partial}\, \frac{\partial \log a_{\widetilde{U}}}{\partial \tilde{z}_k}, \ \ \overline{\partial}\psi_k = 0,$$

Consider the matrix of C^∞ $(1,1)$-forms

$$\widetilde{\Omega} = (\widetilde{\Omega}_{ij}),\ \widetilde{\Omega}_{ij} = \Omega_{ij} - d\tilde{z}_j \wedge \psi_i.$$

In [4] it is shown that, for any invariant polynomial of degree n, whenever $\widetilde{U}_\alpha \cap \widetilde{U}_\beta \neq \emptyset$ the matrices $\widetilde{\Omega}_\alpha$ satisfy $P(\widetilde{\Omega}_\alpha) = P(\widetilde{\Omega}_\beta)$ and hence they define *a global smooth* (n,n)*-form* $P(\widetilde{\Omega})$ *on* X.

Outside the zeros of the section ξ, the form $P(\widetilde{\Omega})$ satisfies

$$P(\widetilde{\Omega}) = d\Psi \tag{10}$$

where Ψ is a smooth $(n, n-1)$-form on $X \setminus \{p : \xi(p) = 0\}$ and

$$\left[\left(\frac{\sqrt{-1}}{2\pi}\right)^n P(\widetilde{\Omega})\right] = P(TX - L^\vee) \in H^{2n}_{dR}(X, \mathbb{C}). \tag{11}$$

Write

$$\tilde{\xi}_\alpha = \sum_i \tilde{\xi}^\alpha_i \frac{\partial}{\partial \tilde{z}^\alpha_i}, \qquad \tilde{\xi}^\alpha_i \in \Gamma(\widetilde{U}_\alpha, \breve{L}_{|\widetilde{U}_\alpha}) \tag{12}$$

for the local expression of $\tilde{\xi}$ over $\widetilde{U}_\alpha$.

At each p_α which is a zero of ξ, adopt the standard hermitian metrics $h^\alpha_{ij} = \delta_{ij}$ and $H^\alpha = 1$ over U_α and extend these to h and H, in $\mathscr{T}_X$ and $\mathscr{L}$ respectively, by means of a

partition of unity. Then, with the induced metrics over $\tilde{U}_\alpha$, the $(n, n-1)$-form Ψ above has the explicit expression (equation (55) of [4]):

$$\pi_\alpha^*\Psi = (-1)^{n+1}P(J\tilde{\xi}_\alpha)\ \eta_\alpha \wedge\ (\overline{\partial}\,\eta_\alpha)^{n-1} \tag{13}$$

where

$$\eta_\alpha = \frac{\sum\limits_i \overline{\tilde{\xi}_i^\alpha}\ d\tilde{z}_i^\alpha}{\sum\limits_i \overline{\tilde{\xi}_i^\alpha}\ \tilde{\xi}_i^\alpha}$$

is the $(0,1)$-form dual to $\tilde{\xi}_\alpha$.

We now investigate $\eta_\alpha \wedge\ (\overline{\partial}\eta_\alpha)^{n-1}$. We have

(14)

$$\eta_\alpha\wedge(\overline{\partial}\eta_\alpha)^{n-1}=\frac{\upsilon_n}{|\tilde{\xi}_\alpha|^{2n}}\left[d\tilde{z}_1^\alpha\wedge\cdots\wedge d\tilde{z}_n^\alpha\bigwedge\sum_i(-1)^{i-1}\overline{\tilde{\xi}_i^\alpha}\,d\,\overline{\tilde{\xi}_1^\alpha}\wedge\cdots\wedge\widehat{d\,\overline{\tilde{\xi}_i^\alpha}}\wedge\cdots\wedge d\,\overline{\tilde{\xi}_n^\alpha}\right]$$

where $\upsilon_n = (-1)^{\frac{n(n-1)}{2}}(n-1)!$.

Recall that the Bochner-Martinelli kernel in $\mathbb{C}^n \times \mathbb{C}^n$ has the representation

$$\mathsf{K}(w,u) = \frac{(n-1)!}{(2\pi\sqrt{-1})^n|w-u|^{2n}}\ \sum_i(\overline{w_i}-\overline{u_i})\,dw_i\ \wedge\left[\bigwedge_{j\neq i}d\,\overline{w_j}\wedge dw_j\right].$$

A manipulation shows that

$$\mathsf{K}(w,0) = (-1)^{\frac{n(n-1)}{2}}\frac{(n-1)!}{(2\pi\sqrt{-1})^n|w|^{2n}}\sum_i\Theta_i(w)\wedge\vartheta(w)$$

where $\Theta_i(w) = (-1)^{i-1}\overline{w_i}\,d\,\overline{w_1}\wedge\cdots\wedge\widehat{d\,\overline{w_i}}\wedge\cdots\wedge d\,\overline{w_n}$ and $\vartheta(w) = dw_1\wedge\cdots\wedge dw_n$. Set $\ell_n = (-1)^{\frac{n(n-1)}{2}}\,\frac{(n-1)!}{(2\pi\sqrt{-1})^n}$ and put

$$\mathsf{B}(z,\zeta) = \frac{\ell_n}{|z-\zeta|^2}\ \sum_i\ \Theta_i(z-\zeta)\wedge\vartheta(\zeta).$$

This is the same as

$$\mathsf{B}(z,\zeta) = \frac{\ell_n}{|z-\zeta|^2}\ \sum_i(-1)^{i-1}(\overline{z_i}-\overline{\zeta_i})\bigwedge_{j\neq i}(d\overline{z_j}-d\overline{\zeta_j})\wedge d\zeta_1\wedge\cdots\wedge d\zeta_n.$$

Now, it is proven in [7], Lemma in page 651, that

$$\mathrm{Res}_0\left[\frac{g\,dz_1\wedge\cdots\wedge dz_n}{f_1\ldots f_n}\right] = \int\limits_{S_\epsilon^{2n-1}} g\,\Phi^*\mathsf{B} \tag{15}$$

where g and $f = (f_1,\ldots,f_n)$ are holomorphic in a neigborhood of $0\in\mathbb{C}^n$, f has an isolated zero at 0, $\Phi : (\mathbb{C}^n,0)\longrightarrow\mathbb{C}^n\times\mathbb{C}^n$ is the map $\Phi(z) = (z+f(z),z)$ and S_ϵ^{2n-1} is the euclidean sphere of radius ϵ centered at 0, for all sufficiently small $\epsilon > 0$.

By replacing f by ξ_α and considering the map $\Phi : (\widetilde{U}_\alpha, \tilde{p}_\alpha) \longrightarrow \mathbb{C}^n \times \mathbb{C}^n$, $\Phi(\tilde{z}_\alpha) = (\tilde{z}_\alpha + \tilde{\xi}_\alpha(\tilde{z}_\alpha), \tilde{z}_\alpha)$, a straightforward calculation gives

$$\Phi^*\mathsf{B} = \left(\frac{1}{2\pi\sqrt{-1}}\right)^n \eta_\alpha \wedge (\overline{\partial}\eta_\alpha)^{n-1}. \tag{16}$$

By (11) it follows that

$$\int_X P(TX - L^\vee) = \int_X \left[\left(\frac{\sqrt{-1}}{2\pi}\right)^n P(\widetilde{\Omega})\right]. \tag{17}$$

For each zero p_α of ξ_α choose a euclidean ball $B_\epsilon(p_\alpha)$ centered at p_α such that $\overline{B_\epsilon(p_\alpha)} \subset U_\alpha$. By Stokes and (10)

(18)
$$\int_{X\setminus\cup_1^k B_\epsilon(p_\alpha)} P\left(\frac{\sqrt{-1}}{2\pi}\widetilde{\Omega}\right) = \int_{X\setminus\cup_1^k B_\epsilon(p_\alpha)} \left(\frac{\sqrt{-1}}{2\pi}\right)^n d\Psi = \sum_1^k \int_{\partial B_\epsilon(p_\alpha)} -\left(\frac{\sqrt{-1}}{2\pi}\right)^n \Psi.$$

Now,

$$\int_{S_\epsilon^{2n-1}(p_\alpha)} -\left(\frac{\sqrt{-1}}{2\pi}\right)^n \Psi = \frac{1}{\#G_{p_\alpha}} \int_{S_\epsilon^{2n-1}(\tilde{p}_\alpha)} -\left(\frac{\sqrt{-1}}{2\pi}\right)^n \pi_\alpha^*\Psi \tag{19}$$

$$= \frac{1}{\#G_{p_\alpha}} \int_{S_\epsilon^{2n-1}(\tilde{p}_\alpha)} -\left(\frac{\sqrt{-1}}{2\pi}\right)^n (-1)^{n+1} P(J\tilde{\xi}_\alpha)\, \eta_\alpha \wedge (\overline{\partial}\eta_\alpha)^{n-1} \tag{20}$$

$$= \frac{1}{\#G_{p_\alpha}} \int_{S_\epsilon^{2n-1}(\tilde{p}_\alpha)} -\left(\frac{\sqrt{-1}}{2\pi}\right)^n (-1)^{n+1} P(J\tilde{\xi}_\alpha)\, (2\pi\sqrt{-1})^n \Phi^*\mathsf{B} \tag{21}$$

$$= \frac{1}{\#G_{p_\alpha}} \int_{S_\epsilon^{2n-1}(\tilde{p}_\alpha)} P(J\tilde{\xi}_\alpha)\, \Phi^*\mathsf{B} = \frac{1}{\#G_{p_\alpha}} \mathrm{Res}_{\tilde{p}_\alpha}\left[\frac{P(J\tilde{\xi}_\alpha)\, d\tilde{z}_1^\alpha \wedge \cdots \wedge d\tilde{z}_n^\alpha}{\tilde{\xi}_1^\alpha \ldots \tilde{\xi}_n^\alpha}\right] \tag{22}$$

by (15). Hence

(23)
$$\int_{X\setminus\cup_1^k B_\epsilon(p_\alpha)} P\left(\frac{\sqrt{-1}}{2\pi}\widetilde{\Omega}\right) = \sum_{p_\alpha \,|\, \xi(p_\alpha)=0} \frac{1}{\#G_{p_\alpha}} \mathrm{Res}_{\tilde{p}_\alpha}\left[\frac{P(J\tilde{\xi}_\alpha)\, d\tilde{z}_1^\alpha \wedge \cdots \wedge d\tilde{z}_n^\alpha}{\tilde{\xi}_1^\alpha \ldots \tilde{\xi}_n^\alpha}\right].$$

Taking $\epsilon \to 0$ we get, by (17),

$$\int_X P(TX - L^\vee) = \sum_{p \,|\, \xi(p)=0} \frac{1}{\#G_p} \mathrm{Res}_{\tilde{p}}\left[\frac{P(J\tilde{\xi})\, d\tilde{z}_1 \wedge \cdots \wedge d\tilde{z}_n}{\tilde{\xi}_1 \ldots \tilde{\xi}_n}\right] \tag{24}$$

as stated. □

2.2. **Another residue formula.** In this subsection we bring in resolutions and obtain relations between characteristic numbers of the orbifold and of a good resolution. As in **2.1**, let X be a compact complex orbifold of dimension n with only isolated singularities, $\mathscr{L}$ a locally V-free sheaf of rank 1 over X and L the associated line V-bundle.

Suppose ξ is a holomorphic section of $TX \otimes L$ with isolated zeros. The Poincaré-Hopf index of ξ at a zero p was defined by Satake in §3.2 of [11] and is:

$$I_p(\xi) = \frac{1}{\#G_p} I_{\tilde{p}}(\tilde{\xi}) \tag{25}$$

where $\pi_p : (\widetilde{U}, \tilde{p}) \longrightarrow (U, p)$ is a local smoothing covering, $\tilde{\xi} = \pi_p^* \xi$ and $I_{\tilde{p}}(\tilde{\xi})$ is the usual Poincaré-Hopf index. It's well known that, in terms of point residues, using $\det$ to designate the n-th elementary symmetric polynomial C_n,

$$I_{\tilde{p}}(\tilde{\xi}) = \operatorname{Res}_{\tilde{p}} \left[\frac{\det(J\tilde{\xi})\, d\tilde{z}_1 \wedge \cdots \wedge d\tilde{z}_n}{\tilde{\xi}_1 \cdots \tilde{\xi}_n} \right]. \tag{26}$$

Recall from (9) and Remark 1 (iv) that

$$c_n(TX \otimes L) = c_n(TX - L^\vee) = c_n(X) + c_{n-1}(X)c_1(L) + \cdots + (c_1(L))^n. \tag{27}$$

Theorem 2.2. *Let X be a compact complex orbifold of dimension n with only isolated singularities, $\mathscr{L}$ a locally V-free sheaf of rank 1 over X and L the associated line V-bundle. Let $\sigma : \widetilde{X} \longrightarrow X$ be a good resolution of X, $\mathcal{E}$ the exceptional divisor, and let ξ be a holomorphic section of $TX \otimes L$ with isolated zeros. Assume $(\sigma^*\mathscr{L})^{\vee\vee}$ is locally free. Then*

(28)

$$\sum_{p\,|\,\xi(p)=0} I_p(\xi) = \int_{\widetilde{X}} c_n\left((\sigma^*(TX \otimes L))^{\vee\vee}\right) - \sum_{p \in Sing(X)} \int_{\widetilde{X}} c_n\left(p, (\sigma^*(TX \otimes L))^{\vee\vee}\right).$$

In particular, if $\sigma^*(\mathscr{T}_X \otimes \mathscr{L})$ is reflexive, then

(29)

$$\sum_{p \in Sing(X) \cap \{\xi(p)=0\}} I_p(\xi) = \sum_{\tilde{p} \in \mathcal{E}\,|\,\sigma^*\xi(\tilde{p})=0} I_{\tilde{p}}(\sigma^*\xi) - \sum_{p \in Sing(X)} \int_{\widetilde{X}} c_n\left(p, \sigma^*(TX \otimes L)\right).$$

Proof. First we remark that since $\mathscr{L}$ is locally V-free, we may construct a resolution such that $(\sigma^*\mathscr{L})^{\vee\vee}$ is locally free (Lemma 3.7 of [2]). By Proposition 1.8, Theorem 2.1, (25) and (26) we have (28). (29) follows from (28) since $\sigma(\mathcal{E}) = Sing(X)$ and $I_p(\xi) = I_{\tilde{p}}(\sigma^*\xi)$ for all $p \notin Sing(X)$. □

3. Applications

3.1. **One-dimensional foliations on $\mathbb{P}^n_w$.** Here we consider weighted complex projective spaces with only isolated singularities, which we briefly recall.

Let $w_0, ..., w_n$ be positive integers two by two co-primes, set $w := (w_0, ..., w_n)$ and $|w| := w_0 + \cdots + w_n$. Define an action of $\mathbb{C}^*$ in $\mathbb{C}^{n+1} \setminus \{0\}$ by

$$\begin{array}{ccc} \mathbb{C}^* \times \mathbb{C}^{n+1} \setminus \{0\} & \longrightarrow & \mathbb{C}^{n+1} \setminus \{0\} \\ \lambda.(z_0, \ldots, z_n) & \longmapsto & (\lambda^{w_0} z_0, \ldots, \lambda^{w_n} z_n) \end{array} \tag{30}$$

and let $\mathbb{P}^n_w := \mathbb{C}^{n+1} \setminus \{0\}/\sim$. The weights are chosen to be 2 by 2 co-primes in order to assure a finite number of singularities and to give $\mathbb{P}^n_w$ the structure of an effective, abelian, compact orbifold of dimension n. The singular locus is:

$$Sing(\mathbb{P}^n_w) = \{\mathsf{e}_i = [0 : \cdots : \underbrace{1}_{i} : \cdots : 0]_w \ : i = 0, 1, \ldots, n\}. \tag{31}$$

We have the canonical projection

$$\begin{array}{rcl} \pi : \mathbb{C}^{n+1} \setminus \{0\} & \longrightarrow & \mathbb{P}^n_w \\ (z_0, \ldots, z_n) & \longmapsto & [z_0^{w_0} : \cdots : z_n^{w_n}]_w \end{array} \tag{32}$$

and the natural map

$$\begin{array}{rcl} \varphi_w : \mathbb{P}^n & \longrightarrow & \mathbb{P}^n_w \\ [z_0 : \cdots : z_n] & \longmapsto & [z_0^{w_0} : \cdots : z_n^{w_n}]_w \end{array} \tag{33}$$

of degree $\deg \varphi_w = w_0 \ldots w_n$. The map φ_w is *good* in the sense of [5] §4.4, which means, among other things, that V-bundles behave well under pullback. It is shown in [8] that there is a line V-bundle $\mathscr{O}_{\mathbb{P}^n_w}(1)$ on $\mathbb{P}^n_w$, unique up to isomorphism, such that

$$\varphi_w^* \mathscr{O}_{\mathbb{P}^n_w}(1) \cong \mathscr{O}_{\mathbb{P}^n}(1) \tag{34}$$

and, by naturality, $c_1(\varphi_w^* \mathscr{O}_{\mathbb{P}^n_w}(1)) = c_1(\mathscr{O}_{\mathbb{P}^n}(1)) = \varphi_w^* c_1(\mathscr{O}_{\mathbb{P}^n_w}(1))$, from which we obtain the Chern number

$$[\mathbb{P}^n_w] \frown \left(c_1(\mathscr{O}_{\mathbb{P}^n_w}(1))\right)^n = \int\limits_{\mathbb{P}^n_w} \left(c_1(\mathscr{O}_{\mathbb{P}^n_w}(1))\right)^n = \frac{1}{w_0 \ldots w_n} \tag{35}$$

since

$$1 = \int\limits_{\mathbb{P}^n} (c_1(\mathscr{O}_{\mathbb{P}^n}(1)))^n = \int\limits_{\mathbb{P}^n} \varphi_w^* \left(c_1(\mathscr{O}_{\mathbb{P}^n_w}(1))\right)^n = (\deg \varphi_w) \int\limits_{\mathbb{P}^n_w} \left(c_1(\mathscr{O}_{\mathbb{P}^n_w}(1))\right)^n. \tag{36}$$

As usual we set $\mathscr{O}_{\mathbb{P}^n_w}(k) := \mathscr{O}_{\mathbb{P}^n_w}(1)^{\otimes k}$ for $k \in \mathbb{Z}$.

The Euler sequence on $\mathbb{P}^n_w$ reads

$$0 \longrightarrow \underline{\mathbb{C}} \longrightarrow \bigoplus_{i=0}^{n} \mathscr{O}_{\mathbb{P}^n_w}(w_i) \longrightarrow T\mathbb{P}^n_w \longrightarrow 0 \tag{37}$$

where

(i) $1 \longmapsto (w_0 z_0, ..., w_n z_n)$.

(ii) $(P_0, ..., P_n) \longmapsto \pi_* \left(\sum_{i=0}^n P_i \frac{\partial}{\partial z_i}\right)$.

Twisting by $\mathscr{O}_{\mathbb{P}^n_w}(d-1)$ we get

$$0 \longrightarrow \mathscr{O}_{\mathbb{P}^n_w}(d-1) \longrightarrow \bigoplus_{i=0}^{n} \mathscr{O}_{\mathbb{P}^n_w}(w_i + d - 1) \longrightarrow T\mathbb{P}^n_w \otimes \mathscr{O}_{\mathbb{P}^n_w}(d-1) \longrightarrow 0. \tag{38}$$

Lemma 3.1. *If* $d > 1 - \max\limits_{i \neq j}\{w_i + w_j\}$, *then* $H^0(\mathbb{P}^n_w, T\mathbb{P}^n_w \otimes \mathcal{O}_w(d-1)) \neq 0$.

Proof. The pairing $\Omega^r_{\mathbb{P}^n_w} \times \Omega^{n-r}_{\mathbb{P}^n_w} \longrightarrow \Omega^n_{\mathbb{P}^n_w}$ induces $\Omega^r_{\mathbb{P}^n_w} \cong \mathrm{Hom}_{\mathbb{C}}\left(\Omega^{n-r}_{\mathbb{P}^n_w}, \Omega^n_{\mathbb{P}^n_w}\right)$. On the other hand, $K_{\mathbb{P}^n_w} = \mathscr{O}_{\mathbb{P}^n_w}(-|w|)$ and hence $T\mathbb{P}^n_w = (\Omega^1_{\mathbb{P}^n_w})^\vee = \Omega^{n-1}_{\mathbb{P}^n_w} \otimes (\Omega^n_{\mathbb{P}^n_w})^\vee = \Omega^{n-1}_{\mathbb{P}^n_w}(|w|)$. Since $H^0(\mathbb{P}^n_w, T\mathbb{P}^n_w \otimes \mathscr{O}_{\mathbb{P}^n_w}(d-1)) \cong H^0(\mathbb{P}^n_w, \Omega^{n-1}_{\mathbb{P}^n_w}(|w|+d-1))$, the result follows from Corollary 2.3.4 of [6]. □

Hence, if X is a quasi-homogeneous vector field of type w and degree d in $\mathbb{C}^{n+1}$, as in Lemma 3.1, that is, writing $\mathsf{X} = \sum_{i=0}^{n} P_i(z)\frac{\partial}{\partial z_i}$ we have $P_i(\lambda^{w_0}z_0, \ldots, \lambda^{w_n}z_n) = \lambda^{d+w_i-1}P_i(z_0, \ldots, z_n)$, then X defines a one-dimensional holomorphic foliation $\mathcal{F}$ on $\mathbb{P}^n_w$ and, due to (38), $g\mathsf{R}_w + \mathsf{X}$ defines the same foliation, where $\mathsf{R}_w = \sum_0^n w_i z_i \frac{\partial}{\partial z_i}$ is the weighted radial vector field and g is a quasi-homogeneous polynomial of type w and degree d. The integer d is, by definition, the *degree* of the foliation $\mathcal{F}$.

Dually, as $H^0(\mathbb{P}^n_w, T\mathbb{P}^n_w \otimes \mathscr{O}_{\mathbb{P}^n_w}(d-1)) \cong H^0(\mathbb{P}^n_w, \Omega^{n-1}_{\mathbb{P}^n_w}(|w|+d-1))$ we have that a one-dimensional holomorphic foliation on $\mathbb{P}^n_w$ is induced by a quasi-homogeneous $(n-1)$-form $\eta = \sum_0^n A_i dz_1 \wedge \cdots \wedge \widehat{dz_i} \wedge \cdots \wedge dz_n$, where A_i is quasi-homoegenous of degree $d + |w| - w_i - 1$, subjected to the condition $\sum_0^n w_i z_i A_i \equiv 0$.

We now give an enumerative result concerning foliations on $\mathbb{P}^n_w$ and examine some consequences of it.

Proposition 3.2. *Let ξ be a holomorphic section of $T\mathbb{P}^n_w \otimes \mathscr{O}_{\mathbb{P}^n_w}(d-1)$ with only isolated zeros. Then*

$$\sum_{p|\xi(p)=0} \mathcal{I}_p(\xi) = \frac{1}{w_0 \ldots w_n} \sum_{j=0}^{n} C_j(w)(d-1)^{n-j}, \tag{39}$$

where $C_j(w)$ is the j-th elementary symmetric function of the variables $w_0, \ldots, w_n$.

Proof. By Theorem 2.1, (25), (26) and (27) we have that

$$\sum_{p|\xi(p)=0} \mathcal{I}_p(\xi) = \int_{\mathbb{P}^n_w} c_n(T\mathbb{P}^n_w \otimes \mathscr{O}_{\mathbb{P}^n_w}(d-1)). \tag{40}$$

To calculate this Chern number we use the map φ_w. By (27),

$$\begin{aligned} &c_n(T\mathbb{P}^n_w \otimes \mathscr{O}_{\mathbb{P}^n_w}(d-1)) = \\ &= c_n(T\mathbb{P}^n_w) + c_{n-1}(T\mathbb{P}^n_w)c_1(\mathscr{O}_{\mathbb{P}^n_w}(d-1)) + \cdots + (c_1(\mathscr{O}_{\mathbb{P}^n_w}(d-1)))^n. \end{aligned} \tag{41}$$

By (37) and Remark 1 (iii),

$$c(T\mathbb{P}^n_w) = c\left(\bigoplus_{i=0}^{n} \mathscr{O}_{\mathbb{P}^n_w}(w_i)\right). \tag{42}$$

By naturality

$$\varphi_w^*\left(c\left(\bigoplus_{i=0}^{n} \mathscr{O}_{\mathbb{P}^n_w}(w_i)\right)\right) = c\left(\bigoplus_{i=0}^{n} \varphi^*\mathscr{O}_{\mathbb{P}^n_w}(w_i)\right) = c\left(\bigoplus_{i=0}^{n} \mathscr{O}_{\mathbb{P}^n}(w_i)\right) \tag{43}$$

and

$$c\left(\bigoplus_{i=0}^{n}\mathscr{O}_{\mathbb{P}^n}(w_i)\right)=\prod_{i=0}^{n}(1+w_i\mathsf{h})=1+\sum_{i=1}^{n}C_i(w)\mathsf{h}^i \tag{44}$$

where h is the hyperplane class in $\mathbb{P}^n$. Hence, from (41) we get

$$\begin{aligned}&\varphi_w^* c_n(T\mathbb{P}^n_w\otimes\mathscr{O}_{\mathbb{P}^n_w}(d-1))=\\&=\left[C_n(w)+C_{n-1}(w)(d-1)+\cdots+C_1(w)(d-1)^{n-1}+(d-1)^n\right]\mathsf{h}^n.\end{aligned} \tag{45}$$

Integration gives:

$$\begin{aligned}\sum_{i=0}^{n}C_i(w)(d-1)^{n-i} &= \int\limits_{\mathbb{P}^n}\varphi_w^* c_n(T\mathbb{P}^n_w\otimes\mathscr{O}_{\mathbb{P}^n_w}(d-1))\\ &= (\deg\varphi_w)\int\limits_{\mathbb{P}^n_w}c_n(T\mathbb{P}^n_w\otimes\mathscr{O}_{\mathbb{P}^n_w}(d-1))\end{aligned} \tag{46}$$

and the result follows. □

Definition 3.3. *Let ξ be a holomorphic section of $T\mathbb{P}^n_w\otimes\mathscr{O}_{\mathbb{P}^n_w}(d-1)$ and let p be an isolated zero of ξ. p is non-degenerate if, given a local smoothing covering $\pi:(\widetilde{U},\tilde{p})\longrightarrow(U,p)$, the lifted vector field $\tilde{\xi}=\pi^*\xi$ has a non-degenerate singularity at $\tilde{p}$, that is, $\tilde{I}_{\tilde{p}}(\tilde{\xi})=1$.*

Given $\mathsf{e}_j\in Sing(\mathbb{P}^n_w)$ (recall (31)), the local fundamental group G_{e_j} of $\mathbb{P}^n_w$ at e_j is isomorphic to $\boldsymbol{\zeta}_{w_j}$, the group of the w_j-th roots of unity. As an immediate consequence of Proposition 3.2 we have

Corollary 3.4. *Let ξ be a holomorphic section of $T\mathbb{P}^n_w\otimes\mathscr{O}_{\mathbb{P}^n_w}(d-1)$ with only isolated and non-degenerate zeros. Then*

$$\sum_{p|\xi(p)=0}\frac{w_0\ldots w_n}{\#G_p}=\sum_{j=0}^{n}C_j(w)(d-1)^{n-j}. \tag{47}$$

If $\mathcal{F}$ is the foliation induced by ξ and $Sing(\mathcal{F})\cap Sing(\mathbb{P}^n_w)=\emptyset$, then (47) reads

$$(w_0\ldots w_n)\,\#Sing(\mathcal{F})=\sum_{j=0}^{n}C_j(w)(d-1)^{n-j}. \tag{48}$$

Corollary 3.5. *Let ξ be a holomorphic section of $T\mathbb{P}^n_w\otimes\mathscr{O}_{\mathbb{P}^n_w}(d-1)$ with only isolated zeros and denote by $\mathcal{F}$ the induced foliation. Then,*

(i) *$Sing(\mathcal{F})\neq\emptyset$ whenever $d\geq 1$ and, for $n=2$, whenever $d\geq 0$.*
(ii) *$Sing(\mathcal{F})\neq\emptyset$ in case $d-1\nmid C_n(w)$.*

Proof. (i) holds since the right hand side of (39) is positive for $d\geq 1$. The case $n=2$ is also straightforward. As for (ii), recall Lemma 3.1 and notice that, if $Sing(\mathcal{F})=\emptyset$ then (39) becomes $\sum_{j=0}^{n-1}C_j(w)(d-1)^{n-j}=-C_n(w)$ and, by (i), necessarily $d\leq 0$. □

Remark 2. Let $\mathcal{F}$ be a foliation of degree $d\neq 1$ on $\mathbb{P}^2_w$, $w=(w_0,w_1,w_2)$, with isolated singularities only. If all points in $Sing(\mathcal{F})$ are non-degenerate singularities, then $Sing(\mathcal{F})\neq\{\mathsf{e}_0,\mathsf{e}_1,\mathsf{e}_2\}$. To see this assume $Sing(\mathcal{F})=\{\mathsf{e}_0,\mathsf{e}_1,\mathsf{e}_2\}$. Then (39) reads

$\frac{C_2(w)}{w_0w_1w_2} = \frac{(d-1)^2+(d-1)C_1(w)+C_2(w)}{w_0w_1w_2}$. Since $d \neq 1$ we are left with $d = 1 - C_1(w)$. But this contradicts Lemma 3.1 since $d > 1 - \max_{i\neq j}\{w_i + w_j\} > 1 - C_1(w)$.

For a complex surface $\tilde{X}$ and $\tilde{\mathcal{F}}$ a holomorphic foliation on $\tilde{X}$ with isolated singularities, the Baum-Bott index of $\tilde{\mathcal{F}}$ at $\tilde{p} \in Sing(\tilde{\mathcal{F}})$ is defined by

$$BB_{\tilde{p}}(\tilde{\mathcal{F}}) = \text{Res}_{\tilde{p}}\left[\frac{\left(\text{tr}(J\tilde{\xi})\right)^2}{\tilde{\xi}_1\,\tilde{\xi}_2}d\tilde{z}_1\wedge d\tilde{z}_2\right],$$

where $\tilde{\xi} = \tilde{\xi}_1\frac{\partial}{\partial\tilde{z}_1} + \tilde{\xi}_2\frac{\partial}{\partial\tilde{z}_2}$ is a vector field defining $\tilde{\mathcal{F}}$ in a neighborhood of $\tilde{p}$. As was done for the Poincaré-Hopf index, we define the Baum-Bott index of a foliation $\mathcal{F}$ on $\mathbb{P}^2_w$ at an isolated singularity p by

$$BB_p(\mathcal{F}) = \frac{1}{\#G_p}\text{Res}_{\tilde{p}}\left[\frac{\left(\text{tr}(J\tilde{\xi})\right)^2}{\tilde{\xi}_1\,\tilde{\xi}_2}d\tilde{z}_1\wedge d\tilde{z}_2\right] \tag{49}$$

where $\tilde{\xi} = \pi_p^*\xi$, for $\xi \in H^0(T\mathbb{P}^2_w \otimes \mathscr{O}_{\mathbb{P}^2_w}(d-1))$ defining $\mathcal{F}$ and $\pi_p : (\widetilde{U},\tilde{p}) \longrightarrow (U,p)$ a local smoothing covering, with U a good representative of $\{p\}$.

Proposition 3.6. *Let ξ be a holomorphic section of $T\mathbb{P}^2_w \otimes \mathscr{O}_{\mathbb{P}^2_w}(d-1)$ with isolated zeros only. Then*

$$\sum_{p|\xi(p)=0} BB_p(\mathcal{F}) = \frac{1}{w_0w_1w_2}(d+|w|-1)^2. \tag{50}$$

Proof. Since tr denotes the invariant symmetric polynomial C_1, Theorem 2.1 gives

$$\sum_{p|\xi(p)=0} BB_p(\mathcal{F}) = \int_{\mathbb{P}^2_w} c_1^2(T\mathbb{P}^2_w - \mathscr{O}_{\mathbb{P}^2_w}(1-d)) \tag{51}$$

By (9) we have

$$\begin{aligned}
\int_{\mathbb{P}^2_w} c_1^2(T\mathbb{P}^2_w - \mathscr{O}_{\mathbb{P}^2_w}(1-d)) &= \int_{\mathbb{P}^2_w}\left(c_1(T\mathbb{P}^2_w) + c_1(\mathscr{O}_{\mathbb{P}^2_w}(d-1))\right)^2 \\
\text{by (42)} \quad &= \int_{\mathbb{P}^2_w}\left(|w|c_1(\mathscr{O}_{\mathbb{P}^2_w}(1)) + (d-1)c_1(\mathscr{O}_{\mathbb{P}^2_w}(1))\right)^2 \\
&= \frac{1}{\deg\varphi_w}\int_{\mathbb{P}^2}\varphi_w^*\big((d+|w|-1)^2(c_1(\mathscr{O}_{\mathbb{P}^2}(1)))^2\big) \\
&= \frac{1}{w_0w_1w_2}(d+|w|-1)^2\int_{\mathbb{P}^2}(c_1(\mathscr{O}_{\mathbb{P}^2}(1)))^2 \\
&= \frac{1}{w_0w_1w_2}(d+|w|-1)^2.
\end{aligned}$$

□

Let $\mathcal{F}$ be a foliation on $\mathbb{P}^2_w$ and suppose p is an isolated singularity of $\mathcal{F}$. We say that p is *radial* provided the lifting $\tilde{\xi} = \pi_p^*\xi$, of a vector field defining $\mathcal{F}$ around p has radial

linear part at $\tilde{p}$, that is, $\tilde{\xi} = \tilde{z}_1\dfrac{\partial}{\partial \tilde{z}_1} + \tilde{z}_2\dfrac{\partial}{\partial \tilde{z}_2} + \cdots$. Notice that, for a radial singularity, since $J(\tilde{\xi}) = I$ we have $\det J(\tilde{\xi}) = 1$ and $\mathrm{tr}J(\tilde{\xi}) = 2$. It follows that

$$I_p(\xi) = \frac{1}{\#G_p} \text{ and } BB_p(\mathcal{F}) = \frac{4}{\#G_p}. \tag{52}$$

Corollary 3.7. *Let $\mathcal{F}$ be a holomorphic foliation of degree d on $\mathbb{P}^2_w$ with isolated singularities only. If all the singularities of $\mathcal{F}$ are radial, then*

$$d = \frac{1}{3}\left(3 - C_1(w) \pm 2\sqrt{C_1(w)^2 - 3C_2(w)}\right). \tag{53}$$

In particular, on $\mathbb{P}(1,1,k)$ with $k \geq 1$, we have $d = \frac{k-1}{3}$ or $d = 1-k$.

Proof. By Propositions 3.2, 3.6 and by (52):

$$\begin{aligned}\frac{1}{w_0w_1w_2}(d+|w|-1)^2 &= \sum_{p|\xi(p)=0} BB_p(\mathcal{F}) \\ &= 4\sum_{p|\xi(p)=0} \mathcal{I}_p(\xi) \\ &= 4\,\frac{(d-1)^2 + C_1(w)(d-1) + C_2(w)}{w_0w_1w_2}\end{aligned}$$

Noticing that $|w| = C_1(w)$ (53) follows. □

Let's comment on the case $\mathbb{P}(1,1,k)$ in the above corollary. If $k = 1$ then $d = 0$ and, up to automorphism, we have only the foliation on $\mathbb{P}^2$ given by $\omega = z_1dz_0 - z_0dz_1$, which is a pencil. For $k > 1$, up to automorphism, we also have only one foliation of degree $1-k$, analogously the pencil defined by $\omega = z_1dz_0 - z_0dz_1$. Hence, if $k > 1$ and $3 \nmid (k-1)$ then radial foliations are pencils.

3.2. **Hirzebruch surfaces as good resolutions of $\mathbb{P}(1,1,k)$.** The k-th Hirzebruch surface Σ_k, $k \geq 0$, is defined as $\mathbb{P}(\mathscr{O}_{\mathbb{P}^1} \oplus \mathscr{O}_{\mathbb{P}^1}(k))$. These are $\mathbb{P}^1$ bundles over $\mathbb{P}^1$, hence rational ruled surfaces and they have just one ruling, except for $k = 0$ since $\Sigma_0 = \mathbb{P}^1 \times \mathbb{P}^1$. J. Matsuzawa [9] gave the following convenient realization of Σ_k:

$$\Sigma_k = \{([z_0 : z_1 : z_2], [s : t]) \in \mathbb{P}^2 \times \mathbb{P}^1 : s^k z_0 = t^k z_1\}. \tag{54}$$

The projection onto the second factor $\mathrm{pr}_2 : \Sigma_n \longrightarrow \mathbb{P}^1$ gives the ruling and, for $k \geq 1$,

$$D = \{([z_0 : z_1 : z_2], [s : t]) \in \Sigma_k : z_0 = z_1 = 0\} \tag{55}$$

is the unique section with $D \cdot D = -k$. Also, $\mathrm{Pic}(\Sigma_k) = \mathbb{Z}D \oplus \mathbb{Z}L$ where L is a fiber, $D \cdot L = 1$ and $L \cdot L = 0$.

Consider the orbifold $\mathbb{P}(1,1,k)$, $k \geq 1$, which we denote by $\mathbb{P}^2_k$ and recall that, for $k > 1$, $Sing(\mathbb{P}^2_k) = \{\mathsf{e}_2 = [0:0:1]_k\}$. Let

$$\mathbb{V}_k = \{([z_0 : z_1 : z_2]_k, [s : t]) \in \mathbb{P}^2_k \times \mathbb{P}^1 \ : \ sz_0 = tz_1\}. \tag{56}$$

In view of (54), the map

$$\begin{array}{ccc} \mathbb{P}^2 \times \mathbb{P}^1 & \xrightarrow{\Phi} & \mathbb{P}^2_k \times \mathbb{P}^1 \\ ([z_0 : z_1 : z_2], [s : t]) & \longmapsto & ([z_0^{1/k} : z_1^{1/k} : z_2]_k, [s : t]) \end{array} \tag{57}$$

satisfies

$$\Phi_{|\Sigma_k\setminus D} : \Sigma_k \setminus D \longrightarrow \mathbb{V}_k \setminus \{\mathsf{e}_2\} \tag{58}$$

is an isomorphism, in the sense of Definition 1.1. The composite morphism

$$\sigma = \pi_1 \circ \Phi_{|\Sigma_k} : \Sigma_k \longrightarrow \mathbb{P}^2_k \tag{59}$$

where $\pi_1 : \mathbb{P}^2_k \times \mathbb{P}^1 \longrightarrow \mathbb{P}^2_k$ is the projection onto the first factor, gives a good resolution of $\mathbb{P}^2_k$ with exceptional divisor D.

Proposition 3.8. *Let $\xi \in H^0(\mathbb{P}^2_k, T\mathbb{P}^2_k \otimes \mathscr{O}_{\mathbb{P}^2_k}(d-1))$ be a section with isolated zeros only. Write $\sigma^*\xi = \tilde{\xi}$. Then*

$$\sum_{\tilde{p}\in D|\tilde{\xi}=0} I_{\tilde{p}}(\tilde{\xi}) - I_{\mathsf{e}_2}(\xi) = (d^2+kd+k)\left(k-\frac{1}{k}\right). \tag{60}$$

Proof. From (29) of Theorem 2.2 we have

$$I_{\mathsf{e}_2}(\xi) = \sum_{\tilde{p}\in D|\tilde{\xi}=0} I_{\tilde{p}}(\tilde{\xi}) - \int_{\Sigma_k} c_2\left(\mathsf{e}_2, (\sigma^*(T\mathbb{P}^2_k \otimes \mathscr{O}_{\mathbb{P}^2_k}(d-1)))\right) \tag{61}$$

and hence we must calculate $\displaystyle\int_{\Sigma_k} c_2\left(\mathsf{e}_2, (\sigma^*(T\mathbb{P}^2_k \otimes \mathscr{O}_{\mathbb{P}^2_k}(d-1)))\right)$. From (28) and Proposition (3.2) we have, recalling that $w=(1,1,k)$,

(62)

$$\begin{aligned}
&\int_{\Sigma_k} c_2\left(\mathsf{e}_2, (\sigma^*(T\mathbb{P}^2_k \otimes \mathscr{O}_{\mathbb{P}^2_k}(d-1)))\right)\\
&= \int_{\Sigma_k} c_2\left(\sigma^*(T\mathbb{P}^2_k \otimes \mathscr{O}_{\mathbb{P}^2_k}(d-1))\right) \quad - \quad \frac{(d-1)^2+(d-1)C_1(w)+C_2(w)}{k}\\
&= \int_{\Sigma_k} c_2\left(\sigma^*(T\mathbb{P}^2_k \otimes \mathscr{O}_{\mathbb{P}^2_k}(d-1))\right) \quad - \quad \frac{(d-1)^2+(d-1)(k+2)+(2k+1)}{k}\\
&= \int_{\Sigma_k} c_2\left(\sigma^*(T\mathbb{P}^2_k \otimes \mathscr{O}_{\mathbb{P}^2_k}(d-1))\right) \quad - \quad \frac{d^2+kd+k}{k}.
\end{aligned}$$

Now,

$$\begin{aligned}
c_2(\sigma^*(T\mathbb{P}^2_k \otimes \mathscr{O}_{\mathbb{P}^2_k}(d-1))) &= c_2(\sigma^*T\mathbb{P}^2_k) + c_1(\sigma^*T\mathbb{P}^2_k)c_1(\sigma^*\mathscr{O}_{\mathbb{P}^2_k}(d-1))\\
&+ \left(c_1(\sigma^*\mathscr{O}_{\mathbb{P}^2_k}(d-1))\right)^2
\end{aligned}$$

and by using Euler's sequence (37) we get $c_2(\sigma^*T\mathbb{P}^2_k) = (2k+1)\left(c_1(\sigma^*\mathscr{O}_{\mathbb{P}^2_k}(1))\right)^2$ and $c_1(\sigma^*T\mathbb{P}^2_k) = (k+2)c_1(\sigma^*\mathscr{O}_{\mathbb{P}^2_k}(1))$, from which we deduce

$$c_2(\sigma^*(T\mathbb{P}^2_k \otimes \mathscr{O}_{\mathbb{P}^2_k}(d-1))) = (d^2+kd+k)\left(c_1(\sigma^*\mathscr{O}_{\mathbb{P}^2_k}(1))\right)^2. \tag{63}$$

By considering the divisor $S=\{z_0=0\}$ in $\mathbb{P}^2_k$ and noticing that $\mathscr{O}_{\mathbb{P}^2_k}(1) \cong \mathscr{O}_{\mathbb{P}^2_k}(S)$, a straigthforward calculation gives $c_1(\sigma^*\mathscr{O}_{\mathbb{P}^2_k}(1)) = kL + D$, L a fiber, $L\cdot D = 1$, $L^2 = 0$ and hence $\displaystyle\int_{\Sigma_k}\left(c_1(\sigma^*\mathscr{O}_{\mathbb{P}^2_k}(1))\right)^2 = k$. Substituting into (62) we obtain the result. □

The next result gives a condition for a foliation to have a singularity at $Sing(\mathbb{P}^2_k)$. We will give two proofs of it, one as a consequence of Proposition 3.2 and the other by using Proposition 3.8.

Proposition 3.9. *Let $\mathcal{F}$ be a foliation of degree d on $\mathbb{P}^2_k$, $k > 1$. Suppose $\mathcal{F}$ has only isolated singularities. If $k \nmid d^2$ then $\mathcal{F}$ has a singularity at $Sing(\mathbb{P}^2_k) = \{\mathsf{e}_2\}$.*

First proof. In fact, suppose e_2 is non-singular for $\mathcal{F}$. Then the right hand side of (39) is an integer and this happens only if $k \mid d^2$. □

Second proof. If $\mathcal{F}$ is non-singular at e_2, then (60) reads

$$\sum_{\tilde{p}\in D|\tilde{\xi}=0} I_{\tilde{p}}(\tilde{\xi}) = (d^2 + kd + k)\left(k - \frac{1}{k}\right).$$

Since the left hand side is an integer and $k > 1$ we must have $k \mid d^2$. □

Remark 3. Still in $\mathbb{P}^2_k$ we observe that, in case $d \geq 0$ and $\mathsf{e}_2 \in Sing(\mathcal{F})$ is non-degenerate, then $Sing(\mathcal{F})$ contains at least another point. This is because, in case this does not happens, then (39) reads $\dfrac{1}{k} = \dfrac{d^2 + kd + k}{k}$, which is impossible since $k > 1$ and $d \geq 0$.

Example 3.10. Let $\eta = (k+1)f\,dg - g\,df$ where $f(z_0, z_1, z_2) = z_0^{k+1} + z_1^{k+1} - (z_0 + z_1)z_2$ and $g(z_0, z_1, z_2) = az_0^k + bz_1^k + cz_2$, $k > 1$. η induces a foliation $\mathcal{F}$ of degree k on $\mathbb{P}^2_k$, with isolated singularities and $\mathcal{F}$ is non-singular at e_2.

ACKNOWLEGMENTS. The third named author is grateful to IMPA and to the Univ. of Valladolid for hospitality.

REFERENCES

[1] A. ADEM, J. LEIDA, Y. RUAN, *Orbifolds and String Topology*, Cambridge University Press, ISBN-10 0-511-28288-5, 2007.

[2] R. BLACHE, *Chern classes and Hirzebruch-Riemann-Roch theorem for coherent sheaves on complex-projective orbifolds with isolated singularities*, Math. Z., **222** (1996), 7-57.

[3] J-P. BRASSELET, J. SEADE, T. SUWA, *Vector Fields on Singular Varieties*, LNM 1987, Springer-Verlag, Berlin, 2009.

[4] S. S. CHERN, *Meromorphic vector fields and characteristic numbers*, Selected Papers, Springer-Verlag, New York, 1978, 435-443.

[5] W. CHEN AND Y. RUAN, *Orbifold Gromov-Witten theory*, Contemporary Mathematics 310, AMS, 2002, 25-86.

[6] I. DOLGACHEV, *Weighted Projective Varieties*, Lecture Notes in Mathematics 956, Springer-Verlag, 1982, 34-71.

[7] P. GRIFFITHS AND J. HARRIS, *Principles of Algebraic Geometry*, John Wiley & Sons, ISBN 0-471-32792-1, 1978.

[8] É. MANN, *Cohomologie quantique orbifolde des espaces projectifs à poids*, http://arxiv.org/abs/math/0510331v1, 2005.

[9] JUN-ICHI MATSUZAWA *Root Systems and Periods on Hirzebruch Surfaces*, Publ. RIMS, Kyoto Univ. **29** (1993), 411-438.

[10] D. PRILL, *Local classification of quotients of complex manifolds by discontinuous groups*, Duke Math. J. 34 (1967) 375-386.

[11] I. SATAKE, *The Gauss-Bonnet theorem for V-manifolds*, Journal of the Mathematical Society of Japan Vol. 9, No. 4 (1957) 464-492.

[12] W. THURSTON, *The geometry and topology of 3-manifolds*, preprint (1990).

MAURÍCIO CORRÊA JR., DEP. MATEMÁTICA ICEX - UFMG, CAMPUS PAMPULHA, 31270-901 BELO HORIZONTE - BRASIL.
E-mail address: mauricio@mat.ufmg.br

M. RODRIGUEZ PEÑA , DEP. MATEMÁTICA ICEX - UFMG, CAMPUS PAMPULHA, 31270-901 BELO HORIZONTE - BRASIL
E-mail address: miguel.rodriguez.mat@gmail.com

MÁRCIO G. SOARES, DEP. MATEMÁTICA ICEX - UFMG, CAMPUS PAMPULHA 31270-901 BELO HORIZONTE - BRASIL.
E-mail address: msoares@mat.ufmg.br